



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

A Randomized Pilot Study Using Calcitriol in Hospitalized Patients

Yasmine M. Elamir, Hajira Amir, Steven Lim, Yesha Patel Rana, Carolina Gonzalez Lopez, Natalia Viera Feliciano, Ali Omar, William Paul Grist, Michael A. Via



PII: S8756-3282(21)00341-0

DOI: <https://doi.org/10.1016/j.bone.2021.116175>

Reference: BON 116175

To appear in: *Bone*

Received date: 12 June 2021

Revised date: 1 August 2021

Accepted date: 3 September 2021

Please cite this article as: Y.M. Elamir, H. Amir, S. Lim, et al., A Randomized Pilot Study Using Calcitriol in Hospitalized Patients, *Bone* (2021), <https://doi.org/10.1016/j.bone.2021.116175>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

COVER LETTER

To Whom It May Concern,

Thank you for considering our work for potential publication in the prestigious *Bone* Journal

Briefly: The systemic illness associated with SARS-CoV-2 infection results in hospitalization rate of 380.3 hospitalizations per 100,000 population, overwhelming health care systems. Some emerging therapies demonstrate modest benefits, but use is not feasible in large inpatient numbers. Vitamin D regulates expression of approximately 11,000 genes spanning many physiologic functions that include regulation of both innate and adaptive immune function. We investigate the potential benefit of calcitriol therapy given to patients hospitalized with COVID-19. This pilot study is an open label, 1:1 randomized clinical trial of calcitriol 0.5 mcg daily for 14 days or hospital discharge vs. no treatment given to hospitalized adult patients with COVID-19.

We enrolled 50 consecutive patients, 25 per trial arm. The change in peripheral arterial oxygen saturation to the inspired fraction of oxygen (SaO₂/FIO₂ ratio) on admission and discharge between the groups the control group had an average increase of +13.2 (+/- 127.7) on discharge and the calcitriol group had an increase of +91.04 (+/- 112.58) (p= .0305), suggesting an improvement in oxygenation among subjects who received calcitriol. Additionally, 12 patients in the control group required oxygen supplementation on admission and 21 of them were discharged on room air. 14 subjects needed oxygen supplementation in the calcitriol group on admission while all 25 were discharged on room air.

Other clinical markers showed the average length of stay was 9.24 (± 9.4) in the control group compared to 5.5 (± 3.9) days in the calcitriol group (p= .14). The need for ICU transfer was 8 in the control group and 5 in the calcitriol group. There were 3 deaths and 4 readmissions in the control group and 0 deaths and 2 readmissions in the calcitriol group.

This pilot study illustrates the potential benefit for calcitriol use in the treatment of COVID-19 and suggests the need for a larger randomized clinical trial.

The manuscript including all data, figures, tables, and supplementary materials has not been previously reported or published and has not been nor will be submitted to another journal while under review by the *Bone* Journal

There are many publications cited in our manuscript discussion regarding vitamin D and the treatment of ARDS. One study cited was very similar to ours as it studied Calcifediol in hospitalized patients with COVID-19 that corroborates our findings. Another study by Arnein et al examined the use of high dose cholecalciferol in COVID-19.

We would like constructive and direct feedback with as quick of a response as possible to get our work published.

FACE PAGE

Title: A Randomized Pilot Study Using Calcitriol in Hospitalized Patients

Authors: Yasmine M Elamir, MD¹, Hajira Amir, MD¹, Steven Lim, MD², Yesha Patel Rana, MD², Carolina Gonzalez Lopez, MD², Natalia Viera Feliciano, MD², Ali Omar, MD², William Paul Grist, MD³, and Michael A Via, MD¹

¹Division of Endocrinology Diabetes and Bone Disease, Icahn School of Medicine at Mount Sinai Beth Israel, Mount Sinai Morningside, Mount Sinai West

²Division of Internal Medicine at Mount Sinai Beth Israel

³Division of Pulmonary and Critical Care Medicine at St Joseph's University Medical Center

Running Title: Calcitriol and COVID-19

Corresponding Author: Hajira Amir

Address: 75 West End Avenue, Apt. C16F, New York, NY, 10023

Phone Number: 571-344-2533

Email: hajira.amir@gmail.com

DISCLOSURE PAGE

None of the authors have anything to disclose regarding the below.

1. Service on Board of Directors
2. Consulting (other than Advisory Boards or Board of Directors)
3. Position in a company, employment or executive position in pharmaceutical, medical device, or diagnostic companies; including industry scientists in the bone and mineral field
4. Serving as an expert witness or consultant in litigation for commercial entities
5. Honoraria or royalties for books or publications or for lectures (speaker fees) or participating in a speakers bureau
6. Research grants, direct salary support or other financial support from commercial entities
7. Stock holdings and/or stock options in pharmaceutical, medical device, or diagnostic companies
8. Partnerships, warrants, royalties for inventions (licensing revenues) or other ownership interest
9. No institutional investigation regarding conflict of interest, responsible conduct in research, animal welfare, human subjects, or laboratory safety compliance exists
10. Or any other situation or transaction or formal role where we have interest

Data Availability Statement: *The data that support the findings of this study will be openly available in a public archive.*

AUTHORS CONTRIBUTIONS

Yasmine Elamir: Conceptualization, Methodology, Formal analysis, Investigation, Writing - Original Draft, Writing - Review & Editing. **Hajira Amir:** Conceptualization, Methodology, Formal analysis, Investigation, Writing - Original Draft, Writing - Review & Editing. **Steven Lim:** Formal analysis, Investigation. **Yesha Patel Rana:** Formal analysis, Investigation. **Carolina Gonzalez Lopez:** Formal analysis, Investigation. **Natalia Viera Feliciano:** Formal analysis, Investigation. **Ali Omar:** Formal analysis, Investigation. **William Paul Grist:** Formal analysis, Investigation. **Michael A Via:** Conceptualization, Methodology, Formal analysis, Investigation, Writing - Original Draft, Writing - Review & Editing, Supervision.

Abstract

The systemic illness associated with SARS-CoV-2 infection results in hospitalization rate of 380.3 hospitalizations per 100,000 population, overwhelming health care systems. Vitamin D regulates expression of approximately 11,000 genes spanning many physiologic functions that include regulation of both innate and adaptive immune function. We investigate potential benefit of calcitriol therapy given to patients hospitalized with COVID-19.

This was an open label, randomized clinical trial of calcitriol or no treatment given to hospitalized adult patients with COVID-19. Subjects were randomly assigned treatment with calcitriol 0.5 mcg daily for 14 days or hospital discharge; or no treatment (1:1) at time of enrollment.

We enrolled 50 consecutive patients, 25 per trial arm. The change in peripheral arterial oxygen saturation to the inspired fraction of oxygen (SaO₂/FIO₂ ratio) on admission and discharge between the groups the control group had an average increase of +13.2 (+/- 127.7) on discharge and the calcitriol group had an increase of +91.04 (+/- 119.08) (p = .0305), suggesting an improvement in oxygenation among subjects who received calcitriol. Additionally, 12 patients in the control group required oxygen supplementation on admission and 21 of them were discharged on room air. 14 subjects needed oxygen supplementation in the calcitriol group on admission while all 25 were discharged on room air. Other clinical markers showed the average length of stay was 9.24 (± 9.4) in the control group compared to 5.5 (± 3.9) days in the calcitriol group (p = .14). The need for ICU transfer was 8 in the control group and 5 in the calcitriol group. There were 3 deaths and 4 readmissions in the control group and 0 deaths and 2 readmissions in the calcitriol group.

This pilot study illustrates improvement in oxygenation among hospitalized patients with COVID-19 treated with calcitriol and suggests the need for a larger randomized trial.

Key Words: Calcitriol, Vitamin D, COVID-19, Endocrine, Therapy

Introduction

The systemic illness that manifests SARS-CoV-2 infection has resulted in a cumulative hospitalization rate of 380.3 hospitalizations per 100,000 population according to the Centers for Disease Control (CDC). A triggered cytokine storm commonly results in acute respiratory distress syndrome (ARDS), multiorgan system failure, and is potentially fatal in approximately 2-3% of cases, including over 50% of patients who require mechanical ventilation. (1) Emerging therapies including remdesivir and corticosteroids demonstrate modest benefits.(2)

Vitamin D regulates expression of approximately 11,000 genes spanning many physiologic functions that include regulation of both innate and adaptive immune function, pulmonary cytokine release, and induction of autophagy as an immune defense. (3) Synthesis of 1,25-dihydroxyvitamin D (1,25(OH)₂D) is integral to the normal development of antigen-presenting cells. (4) Moreover, through induction of macrophage expression of cathelicidin and β -Defensin2, 1,25(OH)₂D stimulates chemotaxis of neutrophils, monocytes, macrophages, and T cells, promotes the clearance of respiratory pathogens through apoptosis and autophagy of infected epithelial cells. Vitamin D-induced autophagy modulates the immune response in influenza A infection, and diminishes clinical severity of coronavirus infection. (3)

Hospitalized and severely ill patients exhibit high rates of vitamin D deficiency, potentially impacting these immune pathways. (5-7) Lower serum concentration 25-hydroxyvitamin D is associated with adverse outcomes including increased susceptibility to SARS-CoV-2 infection, and increased severity of symptoms in the course of COVID-19.(8, 9)

Activation of vitamin D-dependent pathways may provide clinical benefit to patients with SARS-CoV-2 infection. As the biochemically active metabolite, 1,25(OH)₂D or calcitriol, is useful over other forms of vitamin D as therapy in patients with hypocalcemia or with secondary hyperparathyroidism due to renal insufficiency. (3) We investigate potential activation of vitamin D-dependent pathways in a trial of calcitriol therapy given to patients who are hospitalized with COVID-19.

Materials and Methods

The study protocol was approved by the Mount Sinai Program for the protection of Human subjects/Institutional Review Board (IRB) for enrollment at all sites involved. Verbal consent was obtained from all patients about the outcomes that would be measured, and potential benefits and side effects with subsequent acceptance recorded in the electronic medical record.

Study Design and Participants

This was an open label, randomized clinical trial investigating the use of calcitriol in hospitalized patients with COVID-19. We planned to enroll 50 consecutive hospitalized adult patients with COVID-19 admitted to Mount Sinai Beth Israel, Mount Sinai Morningside, and Mount Sinai West Hospitals. Patients were excluded if they are admitted directly to the intensive care unit (ICU), if they had any of the following: hypercalcemia and/or hyperphosphatemia on admission blood tests, untreated disorders of calcium metabolism including hyperparathyroidism, hypoparathyroidism, chronic renal insufficiency with glomerular filtration rate < 30 ml/min, or if they are prescribed calcitriol for any reason outside of the study. Patients were also excluded if they declined to participate prior to enrollment. Any subject who withdrew from the trial or who left the hospital against medical advice would be included in intention to treat analysis.

Procedures

Eligible patients were allocated at a 1:1 calcitriol, no calcitriol ratio through electronic randomization on the day of admission. Enrolled subjects were assigned treatment with calcitriol 0.5 mcg daily for 14 days or hospital discharge, whichever came first; or no treatment. The remainder of the patient's care was determined by the primary team and may include treatment with remdesivir (200 mg for one day followed by 100mg for 4 days), dexamethasone (6 mg daily for 10 days), or convalescent plasma, as well as supplemental O2. Medication was supplied by the inpatient pharmacy.

Outcomes

Outcomes of the effectiveness of calcitriol in the treatment of COVID-19 included oxygen requirements, length of hospital stay, need for ICU admission, mortality, and readmission.

Laboratory Analysis and Respiratory Function Test

Clinical samples for SARS-CoV-2 diagnostic testing were obtained for all hospital admissions at each medical center by nasopharyngeal exudate sampling. Procedures for RNA extraction and real-time RT-PCR were undertaken at on-site testing using Roche Cobas 6800 System or Agena Mass ARRAY System.

Hematology analyses included a complete blood count (DYN Sapphire, Siemens Healthineers, Erlangen, Germany) and coagulation studies included d-Dimer (Sta R max, 5070194, 5070195). Biochemical tests including renal function (Abbott Architect Analyzer), liver function (Abbott Architect Analyzer), lactate dehydrogenase (Abbott Architect Analyzer Siemens Healthineers,

Erlangen, Germany), ferritin, and, ESR (Excite 40 Random access ESR Analyzer 8060), C-reactive protein (Abbott Architect Analyzer, Siemens Healthineers, Erlangen, Germany, IL-6 (Siemens Healthineers, Erlangen, Germany) Respiratory function was assessed by peripheral oxygen saturation.

To assess respiratory status and oxygen requirement, the ratio of peripheral arterial oxygen saturation to the inspired fraction of oxygen ($\text{SpO}_2/\text{FiO}_2$) was calculated. The peripheral arterial oxygenation was completed with SpO_2 pulse oximeter. The validity of this measure as a surrogate for the pressure arterial oxygen to the fraction of inspired oxygen ($\text{PaO}_2/\text{FiO}_2$) ratio has been demonstrated in numerous studies (10)

Statistical Analysis

Descriptive statistics were used for demographic, laboratory, and clinical prognostic factors related to COVID-19 for each treatment arm. This included age, race, sex, body mass index, comorbidities, COVID-19 treatments utilized, and baseline IL-6 and procalcitonin levels.

The comparison between groups of quantitative variables were performed by using *t*-test for qualitative variables, χ^2 tests and Fisher exact tests (with frequencies <5) were used as well as a Mann-Whitney test for length of stay. The comparison for the improvement in $\text{SpO}_2/\text{FiO}_2$ ratio was performed using *t*-test.

Univariate and multivariate logistic regressions were used to estimate odds ratio and 95 % CIs for the length of stay, rate of readmission, probability of admission to ICU, and mortality rate. Significant p-value was considered when $p < .05$. All the analysis has been done using IBM SPSS Statistics software (SPSS).

Results


We enrolled 50 consecutive patients, 25 per trial arm between September 2020-December 2020. Baseline characteristics of each group is given in table 1. No subject withdrew from the study. Primary outcome and other clinical outcomes are given in figure 1 and table 2. The average length of stay was 9.24 (\pm 9.4) in the control group compared to 5.5 (\pm 3.9) days in the calcitriol group ($p= .14$). The need for transfer to ICU was 8 in the control group and 5 in the calcitriol group ($p=.33$). There were 3 deaths and 4 readmissions in the control group and 0 deaths and 2 readmissions in the calcitriol group. 12 patients in the control group required oxygen supplementation on admission and 21 of them were discharged on room air. 14 subjects needed oxygen supplementation in the calcitriol group on admission while all 25 were discharged on room air.

When comparing change in oxygen saturation using SaO₂/FIO₂ ratio on admission and discharge between the groups the control group had an average increase of +13.2 (\pm 127.7) on discharge and the calcitriol group had an increase of +91.04 (\pm 119.08) ($p= .0305$) (figure 1). No adverse effects were observed (table 3), including no hypercalcemia or hyperphosphatemia.

Table 1. Baseline Characteristics

	Control Group	Calcitriol Group	p-value
<u>Demographics</u>	25	25	
Age (y)	64 \pm 16	69 \pm 18	0.16
Above age 65	19	14	0.14
Male Gender	13	12	0.77
Race			0.90
Caucasian	5	7	
Non-White Latino	15	14	
Black	3	2	
Asian	2	2	
<u>Poor Prognostic Factors</u>			
Body Mass Index (kg/m ²)	27.2 \pm 5	27.3 \pm 6	0.48
BMI > 30 kg/m ²	6	4	0.48
Hypertension	17	13	0.24
Type 2 Diabetes	11	9	0.56
Cancer History	1	1	1.00
Asthma/COPD	3	5	0.44
<u>Admission Levels</u>			
Procalcitonin	0.21 \pm 0.27	0.11 \pm 0.1	0.14
IL-6	65 \pm 114	38 \pm 27	0.29

Lymphocytes (Absolute Number)	1.25 ± 0.68	1.14 ± 0.55	0.51
C-Reactive Protein	160.7 ± 381	78.9 ± 76.8	0.33
Ferritin	570 ± 667.2	902.6 ± 981.8	0.18
D-Dimer	1.36 ± 0.91	1.65 ± 2.08	0.55
Glomerular Filtration Rate	75 ± 41	69 ± 23	0.28
Oxygen Requirements on Admission			0.74
Room Air	13	11	
Nasal Canula	8	12	
High Flow Nasal Canula	1	0	
Non-rebreather	3	1	
Bipap	0	1	
<u>Treatments during admission</u>			
Remdesivir	11	5	0.25
Dexamethasone	12	13	0.78
Convalescent Plasma	2	4	0.38
<u>Medical Center</u>			0.67
MS Beth Israel	13	15	
MS Morningside	12	7	
MS West	2	3	

Figure 1. SaO_2/FIO_2 

	Control Group N=25	Calcitriol Group N=25	
$\Delta S/F$ Ratio, Mean \pm SD	13.2 \pm 127.7	91.04 \pm 119.08	P=0.0305

Table 2. Clinical outcomes

	Control	Calcitriol	p value
Length of Hospital Stay (days)	9.24 \pm 9.4	5.5 \pm 3.9	0.14
ICU Admission	8	5	0.33
Endotracheal Intubation	2	0	0.48
Mortality	3	0	0.23
Readmission within 30 days	4	2	0.67

Table 3. Adverse effects

	Control	Calcitriol	p value
Hypercalcemia	0	0	-

Hyperphosphatemia	0	0	-
Renal Calculus	0	0	-
Reduction in glomerular filtration rate by >10%	4	0	0.1

Discussion

Our study shows a significant reduction in oxygen requirements in patients hospitalized with COVID-19 who received calcitriol. Since ICU patients were not included, routine arterial blood gas assessment was not available. The PaO₂/FIO₂ ratio is a highly regarded parameter of respiratory function and serves as a predictor of ICU mortality in patients with sepsis and ARDS. (11) Moreover, SaO₂/FIO₂ is commonly used as an end point in clinical trials for patients with COVID-19, with success and is considered an accurate predictor of PaO₂/FIO₂ ratio. (10) ARDS severity when measure by PaO₂/FiO₂ is classified as mild if 200-300, moderate if 100-200, and severe is less than 100. SaO₂/FIO₂ ratios of 181 and 235 corresponded approximately to PaO₂/FIO₂ ratios of 300 and 200, respectively. (12) While infected with COVID-19, an average increase in SaO₂/FIO₂ of 91.04±119.08, as noted with calcitriol therapy, demonstrates a significant clinical improvement, and may render a typical patient to a more mild case of ARDS. The SaO₂/FIO₂ ratio is a reliable noninvasive surrogate of the PaO₂/FIO₂ ratio with the advantage of replacing invasive arterial blood gases. (12)

Effects of calcitriol therapy on other clinical parameters, including hospital length of stay, need for ICU admission, endotracheal intubation, hospital readmission, and mortality did not show significant benefit. This may be secondary to the small size of the trial. In each case, numerical results may favor calcitriol therapy. A larger trial is warranted to investigate further.

On the basis of improved oxygenation, our results are somewhat consistent with those reported by Castillo, et al., who conducted a similar open-label pilot trial examining the effects of calcifediol in hospitalized patients with COVID-19. (13) A significant reduction in ICU admission is noted with calcifediol therapy, again suggesting potential respiratory benefit of therapy with vitamin D derivatives.

In contrast, results of a recent trial demonstrate no clinical benefit from a single dose of 200,000 IU cholecalciferol in patients with moderate to severe COVID-19. (14) While increased serum levels of 25-hydroxyvitamin D were observed, levels of 1,25(OH)₂D were not reported. The efficiency of renal and other systemic 1 α -hydroxylase activity to generate 1,25-dihydroxyvitamin D during acute illness is unknown. (15) Moreover, reduced circulation of vitamin D binding protein (VDBP) during acute illness may diminish clinical activity of cholecalciferol. (16) Calcitriol, but not cholecalciferol, can restore calcium homeostasis in patients with inherited VDBP deficiency. (17) These potential mechanisms suggest administration of calcitriol may be more effective than cholecalciferol in severe acute illness such as COVID-19.

Several preclinical and clinical studies suggest therapy with vitamin D suggest beneficial in ARDS specifically. Animal models demonstrate vitamin D sufficient state mitigates ARDS that is induced by lipopolysaccharide, compared to vitamin D deficiency. (18) Moreover, animal models of ARDS with knockout mutations of the vitamin D receptor (VDR) show significantly worse lung histology and nearly twice the mortality rate as wild type mice. (19) Results of animal model of ARDS in VDR-knockout mice

that were also given losartan suggests that vitamin D functions, in part, by up-regulating the pulmonary renin/angiotensin system to mitigate lung damage. (19) This same system is disrupted by COVID-19 infection. (20)

Human trials with cholecalciferol therapy for ARDS have mixed results. In a randomized trial, patients that underwent esophagectomy demonstrated a transient reduction in markers of ARDS with 300,000 IU treatment of cholecalciferol compared to placebo. (21) The authors argue that declining levels of VDBP that were observed in this trial may explain the short-lived benefit of cholecalciferol that was observed.

In another randomized trial of 475 mechanically ventilated ICU patients given 540,000 units cholecalciferol vs placebo, a 44% reduction in hospital mortality was observed in the cholecalciferol treated patients among the subgroup of patients with 25-hydroxyvitamin D levels less than 12 ng/ml. (22) In addition, the cholecalciferol treated patients demonstrated a further reduction in mortality that persisted for 6 months of follow up. (22) A larger, multicenter trial is currently underway. (23)

Limitations of our trial include lack of a placebo and lack of blinding. It is unlikely this would affect the SaO₂/FIO₂ endpoint. Additionally, 25-hydroxyvitamin D levels were not measured in our study. Enrollment of patients without regard to vitamin D status strengthens the generalizability of the results.

Sample size was another limitation. The patients in the control group had a higher percentage of patients above 65 years of age and certain comorbidities such as hypertension and diabetes mellitus were higher in that group. The statistical non significance in these baseline characteristics could be due to the small sample size.

The low cost and high availability make calcitriol an attractive candidate for therapy in hospitalized patients with moderate to severe COVID-19. Further study of this approach is warranted.

ACKNOWLEDGEMENTS

There are no acknowledgements.

Works Cited

1. Guan WJ, Zhong NS. Clinical Characteristics of Covid-19 in China. Reply. N Engl J Med. 2020;382(19):1861-2.
2. Grein J, Ohmagari N, Shin D, Diaz G, Asperges E, Castagna A, et al. Compassionate Use of Remdesivir for Patients with Severe Covid-19. N Engl J Med. 2020;382(24):2327-36.
3. Bilezikian JP, Bikle D, Hewison M, Lazaretti-Castro M, Fehrentz AM, Gupta A, et al. MECHANISMS IN ENDOCRINOLOGY: Vitamin D and COVID-19. Eur J Endocrinol. 2020;183(5):R133-R47.
4. Hewison M, Freeman L, Hughes SV, Evans KN, Bland R, Eliopoulos AG, et al. Differential regulation of vitamin D receptor and its ligand in human monocyte-derived dendritic cells. J Immunol. 2003;170(11):5382-90.
5. Nierman DM, Mechanick JL. Bone hyperresorption is prevalent in chronically critically ill patients. Chest. 1998;114(4):1122-8.
6. Van den Berghe G, Van Roosbroeck P, Vanhove P, Wouters PJ, De Pourcq L, Bouillon R. Bone turnover in prolonged critical illness: effect of vitamin D. J Clin Endocrinol Metab. 2003;88(10):4623-32.
7. Lee P, Eisman JA, Center JR. Vitamin D deficiency in critically ill patients. N Engl J Med. 2009;360(18):1912-4.
8. Laird E, Rhodes J, Kenny RA. Vitamin D and Inflammation: Potential Implications for Severity of Covid-19. Ir Med J. 2020;113(5):81.
9. Rhodes J, Dunstan F, Laird E, Subramanian S, Kenny RA. COVID-19 mortality increases with northerly latitude after adjustment for age suggesting a link with ultraviolet and vitamin D. BMJ Nutr Prev Health. 2020;3(1):118-20.
10. Catoire P, Tellier F, de la Riviere C, Beauvieux MC, Valdenaire G, Galinski M, et al. Assessment of the SpO₂/FiO₂ ratio as a tool for hypoxemia screening in the emergency department. Am J Emerg Med. 2021;44:116-20.
11. Santana AR, de Sousa JL, Amorim FF, Menezes BM, Araújo FVB, Soares FB, et al. SaO₂/FiO₂ ratio as risk stratification for patients with sepsis. Critical Care. 2013;17(4):P51.
12. Bilan N, Dastranji A, Ghalehgalab Behbahani A. Comparison of the spo₂/fio₂ ratio and the pao₂/fio₂ ratio in patients with acute lung injury or acute respiratory distress syndrome. J Cardiovasc Thorac Res. 2015;7(1):28-31.
13. Entrenas Castillo M, Entrenas Costa LM, Vaquero Barrios JM, Alcala Diaz JF, Lopez Miranda J, Bouillon R, et al. "Effect of calcifediol treatment and best available therapy versus best available therapy on intensive care unit admission and mortality among patients hospitalized for COVID-19: A pilot randomized clinical study". J Steroid Biochem Mol Biol. 2020;203:105751.

14. Murai IH, Fernandes AL, Sales LP, Pinto AJ, Goessler KF, Duran CSC, et al. Effect of a Single High Dose of Vitamin D3 on Hospital Length of Stay in Patients With Moderate to Severe COVID-19: A Randomized Clinical Trial. *JAMA*. 2021;325(11):1053-60.
15. Zehnder D, Bland R, Williams MC, McNinch RW, Howie AJ, Stewart PM, et al. Extrarenal expression of 25-hydroxyvitamin d(3)-1 alpha-hydroxylase. *J Clin Endocrinol Metab*. 2001;86(2):888-94.
16. Waldron JL, Ashby HL, Cornes MP, Bechervaise J, Razavi C, Thomas OL, et al. Vitamin D: a negative acute phase reactant. *J Clin Pathol*. 2013;66(7):620-2.
17. Henderson CM, Fink SL, Bassyouni H, Argiropoulos B, Brown L, Laha TJ, et al. Vitamin D-Binding Protein Deficiency and Homozygous Deletion of the GC Gene. *N Engl J Med*. 2019;380(12):1150-7.
18. Dancer RC, Parekh D, Lax S, D'Souza V, Zheng S, Bassford CR, et al. Vitamin D deficiency contributes directly to the acute respiratory distress syndrome (ARDS). *Thorax*. 2015;70(7):617-24.
19. Kong J, Zhu X, Shi Y, Liu T, Chen Y, Bhan I, et al. VDR attenuates acute lung injury by blocking Ang-2-Tie-2 pathway and renin-angiotensin system. *Mol Endocrinol*. 2013;27(12):2116-25.
20. Vaduganathan M, Vardeny O, Michel T, McMurray JJV, Pfeffer MA, Solomon SD. Renin-Angiotensin-Aldosterone System Inhibitors in Patients with Covid-19. *N Engl J Med*. 2020;382(17):1653-9.
21. Parekh D, Dancer RCA, Scott A, D'Souza VK, Howells PA, Mahida RY, et al. Vitamin D to Prevent Lung Injury Following Esophagectomy-A Randomized, Placebo-Controlled Trial. *Crit Care Med*. 2018;46(12):e1128-e35.
22. Amrein K, Schnedl C, Holl A, Riedl R, Christopher KB, Pachler C, et al. Effect of high-dose vitamin D3 on hospital length of stay in critically ill patients with vitamin D deficiency: the VITdAL-ICU randomized clinical trial. *JAMA*. 2014;312(15):1520-30.
23. Amrein K, Parekh D, Westrich S, Preiser JC, Berghold A, Riedl R, et al. Effect of high-dose vitamin D3 on 28-day mortality in adult critically ill patients with severe vitamin D deficiency: a study protocol of a multicentre, placebo-controlled double-blind phase III RCT (the VITDALIZE study). *BMJ Open*. 2019;9(11):e031083.

Highlights

Vitamin D has potent immunomodulatory effects on innate and adaptive immune system

Vitamin D associated actions suppresses acute lung injury

Studies show possible association with Vitamin D deficiency, ARDS and cytokine storm

In our study calcitriol improves oxygenation in patients hospitalized with COVID 19